


AMENDMENTS TO THE CLAIMS

A detailed listing of all claims that are, or were, in the present application, irrespective of whether the claim(s) remains under examination in the application are presented below. The claims are presented in ascending order and each includes one status identifier. Those claims not cancelled or withdrawn but amended by the current amendment utilize the following notations for amendment: 1. deleted matter is shown by strikethrough for six or more characters and double brackets for five or less characters; and 2. added matter is shown by underlining.

1. (Previously Presented) An organophotoreceptor comprising an electrically conductive substrate and a photoconductive element on the electrically conductive substrate, the photoconductive element comprising:


a) a charge transport material having the following formula:



where R comprises a hydrogen, an alkyl group, an alkenyl group, a heterocyclic group, or an aromatic group; X comprises an arylamine group selected from the group consisting of a carbazole group, a julolidine group and a p-N,N-diphenylaminophenylene group; and Y comprises a 9-fluorenylidene group having at least a solubilizing substituent, wherein the solubilizing substituent comprises a $-(\text{CH}_2)_n\text{H}$ group where n is an integer between 1 and 50, and one or more of the methylene groups is optionally replaced by O, S, B, P, C=O, O=S=O, a heterocyclic group, an aromatic group, a CR_cR_d group, or a SiR_eR_f where R_c , R_d , R_e , and R_f are, each independently, H, a hydroxyl group, a thiol group, a carboxyl group, an amino group, an alkyl group, an alkoxy group, an alkenyl group, a heterocyclic group, or an aromatic group, or part of a ring group; and

(b) a charge generating compound.

2. (Previously Presented) An organophotoreceptor according to claim 1 wherein X comprises a carbazole group, or a julolidine group.
3. (Original) An organophotoreceptor according to claim 1 wherein the solubilizing substituent comprises a $-C(=O)O-R_5$ group where R_5 is an alkyl group, an alkenyl group, or an aromatic group.
4. (Original) An organophotoreceptor according to claim 1 wherein the 9-fluorenylidene group further comprises at least a substituent selected from the group consisting of a halogen, a NO_2 group, a cyano group, a hydroxyl group, a thiol group, a carboxyl group, an amine group, an ester group, an alkyl group, an alkoxy group, an alkenyl group, and an aromatic group.
5. (Original) An organophotoreceptor according to claim 1 wherein the photoconductive element further comprises a second charge transport material.
6. (Original) An organophotoconductor according to claim 5 wherein the second charge transport material comprises an electron transport compound.
7. (Original) An organophotoreceptor according to claim 1 wherein the organophotoreceptor is in the form of a drum or a belt.
8. (Previously Presented) An electrophotographic imaging apparatus comprising:
 - (a) a light imaging component; and
 - (b) an organophotoreceptor oriented to receive light from the light imaging component, the organophotoreceptor comprising an electrically conductive substrate and a photoconductive element on the electrically conductive substrate, the photoconductive element comprising:
 - (i) a charge transport material having the formula



where R comprises a hydrogen, an alkyl group, an alkenyl group, a heterocyclic group, or an aromatic group; X comprises an arylamine group selected from the group consisting of a carbazole group, a julolidine group and a p-N,N-diphenylaminophenylene group; and Y comprises a 9-fluorenylidene group having at least a solubilizing substituent, wherein the solubilizing substituent comprises a $-(\text{CH}_2)_n\text{H}$ group where n is an integer between 1 and 50, and one or more of the methylene groups is optionally replaced by O, S, B, P, C=O, O=S=O, a heterocyclic group, an aromatic group, a CR_cR_d group, or a SiR_eR_f where R_c , R_d , R_e , and R_f are, each independently, H, a hydroxyl group, a thiol group, a carboxyl group, an amino group, an alkyl group, an alkoxy group, an alkenyl group, a heterocyclic group, or an aromatic group, or part of a ring group; and

(ii) a charge generating compound.

9. (Previously Presented) An electrophotographic imaging apparatus according to claim 8 wherein X comprises a carbazole group, or a julolidine group.

10. (Original) An electrophotographic imaging apparatus according to claim 8 wherein the solubilizing substituent comprises a $-\text{C}(=\text{O})\text{O}-\text{R}_5$ group where R_5 is an alkyl group, an alkenyl group, or an aromatic group.

11. (Original) An electrophotographic imaging apparatus according to claim 8 wherein the 9-fluorenylidene group further comprises at least a substituent selected from the group consisting of a halogen, a NO_2 group, a cyano group, a hydroxyl group, a thiol group, a carboxyl group, an amine group, an ester group, an alkyl group, an alkoxy group, an alkenyl group, and an aromatic group.


12. (Original) An electrophotographic imaging apparatus according to claim 8 wherein the photoconductive element further comprises a second charge transport material.

13. (Original) An electrophotographic imaging apparatus according to claim 12 wherein the second charge transport material comprises an electron transport compound.

14. (Original) An electrophotographic imaging apparatus according to claim 8 further comprising a toner dispenser.

15-20 (Cancelled)

21. (Previously Presented) A charge transport material having the following formula,




where R comprises a hydrogen, an alkyl group, an alkenyl group, a heterocyclic group, or an aromatic group; X comprises an arylamine group selected from the group consisting of a carbazole group, a julolidine group and a p-N,N-diphenylaminophenylene group; and Y comprises a 9-fluorenylidene group having at least a solubilizing substituent, wherein the solubilizing substituent comprises a $-(\text{CH}_2)_n\text{H}$ group where n is an integer between 1 and 50, and one or more of the methylene groups is optionally replaced by O, S, B, P, C=O, O=S=O, a heterocyclic group, an aromatic group, a CR_cR_d group, or a SiR_eR_f where R_c , R_d , R_e , and R_f are, each independently, H, a hydroxyl group, a thiol group, a carboxyl group, an amino group, an alkyl group, an alkoxy group, an alkenyl group, a heterocyclic group, or an aromatic group, or part of a ring group.

22. (Previously Presented) A charge transport material according to claim 21 wherein X comprises a carbazole group, or a julolidine group.

23. (Original) A charge transport material according to claim 21 wherein the solubilizing substituent comprises a $-\text{C}(=\text{O})\text{O}-\text{R}_5$ group where R_5 is an alkyl group, an alkenyl group, or an aromatic group.

24. (Original) A charge transport material according to claim 21 wherein the 9-fluorenylidene group further comprises at least a substituent selected from the group consisting of a halogen, a NO₂ group, a cyano group, a hydroxyl group, a thiol group, a carboxyl group, an amine group, an ester group, an alkyl group, an alkoxy group, an alkenyl group, and an aromatic group.

25. (Previously Presented) A charge transport material according to claim 21 wherein the charge transport material has formula:



where R₆ comprises a hydrogen, an alkyl group, an alkenyl group, a heterocyclic group, or an aromatic group; R₇ comprises a -(CH₂)_nH group where n is an integer between 1 and 50, and one or more of the methylene groups is optionally replaced by O, S, B, P, C=O, O=S=O, a heterocyclic group, an aromatic group, a CR_cR_d group, or a SiR_eR_f where R_c, R_d, R_e, and R_f are, each independently, H, a hydroxyl group, a thiol group, a carboxyl group, an amino group, an alkyl group, an alkoxy group, an alkenyl group, a heterocyclic group, or an aromatic group, or part of a ring group; R₈ comprises a hydrogen, a halogen, a NO₂ group, a cyano group, a hydroxyl group, a thiol group, a carboxyl group, an amine group, an ester group, an alkyl group, an alkoxy group, an alkenyl group, or an aromatic group; and X comprises an arylamine group selected from the group consisting of a carbazole group, a julolidine group and a p-N,N-diphenylaminophenylene group.

26. (Original) A charge transport material according to claim 25 wherein R₈ is a hydrogen and R₇ comprises a -C(=O)O-R₁₃ group where R₁₃ is an alkyl group, an alkenyl group, or an aromatic group.

27. (Previously Presented) A charge transport material according to claim 25 wherein X comprises a carbazole group, or a julolidine group.